

**Probability distribution of density fluctuations in the non-linear
regime**

Probability distribution of density fluctuations in the
non-linear regime

J. Betancort-Rijo

Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain.

M. López-Corredoira

Instituto de Astrofísica de Canarias, E-38200 La Laguna, Tenerife, Spain.

Astronomisches Institut der Universität Basel. Venusstrasse 7. CH-4102 Binningen,
Switzerland (actual address).

Received _____; accepted _____

ABSTRACT

We present a general procedure for obtaining the present density fluctuation probability distribution given the statistics of the initial conditions. The main difficulties faced with regard to this problem are those related to the non-linear evolution of the density fluctuations and those posed by the fact that the fields we are interested in are the result of filtering an underlying field with structure down to scales much smaller than that of filtering. The solution to the latter problem is discussed here in detail and the solution to the former is taken from a previous work.

We have checked the procedure for values of the rms density fluctuation as large as $3/2$ and several power spectra and found that it leads to results in excellent agreement with those obtained in numerical simulations. We also recover all available exact results from perturbation theory.

Subject headings:

1. Introduction

The current model for the formation of large-scale structure is based on the gravitational instability of small initial fluctuations in the energy density. These are assumed to have originated from quantum fluctuations during the inflationary epoch. The fluctuations form a random field, which is fully specified by the n -point joint probability density functions (hereafter PDFs). The one-point PDF is a useful tool for testing the statistical character of the initial conditions (i.e. whether they are Gaussian or others), since its evolution does not depend on the nature of the dark matter (Trimble 1987). So it would be rather convenient to develop an efficient and accurate procedure for obtaining the present PDFs given that for the initial conditions. Many attempts have been made at solving this problem. Hoffman (1987) used the Zel’dovich (1970) approximation for computing the variance of the density fluctuations in the mildly non-linear regime ($\sigma^2 \leq 1$). However, he did not take into account the fact that evolution and filtering are non-commuting operations. To obtain the variance on comoving scale r , one should evolve the whole field (with all its structure) and then filter it on scale r . But in the analytical procedure followed by Hoffman the field is first filtered and then evolved. In the linear regime ($\sigma \ll 1$) both procedures lead to the same result. However, for $\sigma \sim 1$ the results obtained by these procedures are rather different.

A procedure for obtaining the present PDF given the statistics of the initial conditions, that accounts for non-commutability of evolution and filtering was first described by one of us (Betancort-Rijo 1991). This procedure was the result of taking into account the simple fact that overdense regions with present comoving size r correspond to larger comoving regions in the initial field. However, in developing this idea a couple of subtle points were involved that were not properly dealt within that paper.

During more than a decade that has passed since these first attempts at dealing with the problem under consideration (the article by Betancort-Rijo was in fact written in 1988)

much attention have been given to it. New methods have been developed for approximating the non-linear evolution (Buchert 1989; Bouchet et al. 1995) and for the computation of statistics of the evolved field. Bernardeau (1994a) developed a procedure for taking into account the effect of filtering on the moments of the PDF in the small variance limit and used it to compute the first few hierarchical amplitudes. Then, assuming that these quantities do not depend on σ , he obtained (Bernardeau & Kofman 1995) the present PDF for any value of σ . However, this assumption, although it holds for some initial conditions, is not universal (as we shall see) and, even when it holds, it is not easy to justify a priori.

Despite all the attention that has been devoted to the present problem, the idea described by Betancort-Rijo (1991), which is at the root of the possibility of developing a simple and direct (it is not necessary to obtain the vertex generation function) procedure for computing the present PDF in the non-linear regime analytically, has not been followed. The exception is an article by Padmanabhan & Subramanian (1993), but there the problem is treated in the same way as was by Betancort-Rijo (1991), thereby having the same drawbacks.

In this paper we shall show how to correct the errors included in previous work and give the correct procedure for obtaining the present PDF. To check the correctness of the procedure we compute the values of S_3 and S_4 in the low- σ limit (that have been computed analytically by Bernardeau 1994a) for several power spectra. Our procedure should render the exact values of all hierarchical amplitudes in this limit, and this is what is found for these two amplitudes. We also study the dependence on σ of these amplitudes and of the variance in the low limit and for finite values of σ . We then compute the PDF explicitly for several cases for which numerical simulations are available and find them to agree with the numerical results.

In §3 we briefly review the basic idea (fully described in Betancort-Rijo 1991) involved

in the present approximation and show how to use it to derive correctly the present PDF given the initial one. In the second section we shall discuss the technique that we use for the treatment of the non-linear evolution. In the last section we present and discuss the results of several explicit calculations.

2. Non-linear evolution

The complete Zel’dovich approximation (CZA; Betancort-Rijo & López-Corredoira 2000) is an approximation to the non-linear evolution of density fluctuations depending only on three quantities, λ_i , defined so that $1 - \lambda_i$ are the eigenvalues of the tensor $\frac{\partial \vec{u}}{\partial q}$, where $\vec{u}$ stands for the peculiar velocity of matter as given by the linear theory in comoving units, i.e. the peculiar velocity $\vec{v} = R(t)\vec{u}$; $R(t)$ being the scale factor of the Universe and q stands for the Lagrangian coordinates, which are equal to the initial comoving coordinates (Eulerian). This tensor we call the linear local deformation tensor. The full local deformation tensor is defined in the same manner but using the exact value of $\vec{u}$. This approximation is exact to the second order. An approximation exact to the fourth order may be obtained out of it but this implies introducing additional variables. It takes the form (Betancort-Rijo & López-Corredoira 2000)

$$(1 + \delta)^{-1} = \prod_i (1 - r_i(\lambda)\lambda_i + f_1(\lambda)x_i + f_2(\lambda)u_i), \quad (1)$$

where f_1 , f_2 , r_i are certain functions of the variables λ_i . The variables x , u which are given at any point by certain integrals over the whole field (Betancort-Rijo 2001) but, for statistical purposes we only need their probability distribution for fixed values of the λ_i . In fact, to obtain the moments exactly to the third order and almost exactly to the fourth we can use expression (1) with u equal to zero and x equal to its average value for fixed values of the λ_i ($\langle x_i \rangle(\lambda)$). This is the approximation that in principle we use. It

has a dependence on the power spectra through the mean value of x (see Betancort-Rijo & López-Corredoira 2000). However, the dependence is so small that may be neglected without practical lost of accuracy. So, what we do in practice is to use simply the CZA (expression (1) with $x = u = 0$), which is strictly exact to the third order for a particular power spectra (such that $\langle x_i \rangle (\lambda) = 0$) and practically exact for all of them. The CZA gives the density fluctuation at a point in term of the values of the λ_i at that point. When dealing with filtered fields, however, the relevant quantity is the mean density within certain region (of any prescribed shape). It may be shown that this mean density fluctuation is also given exactly at least to the third order by expression (1) with the λ_i representing now the linear eigenvalues of the deformation tensor corresponding to the filtered field; that is, the deformation tensor derived from the peculiar velocity field corresponding to the growing mode of the filtered field. The x variables are now defined in a different manner. For non-filtered fields the presence of this variables is due to the different rate of growth of the various contributions to the local tidal field. In the present case, however, these variables have an extra component due to the filtering process. But, despite of the different origin of the x variables, expression (1) and the probability distributions for the x retain their form at least up to third order. So, setting in this expression x equal to its mean value for fixed λ_i leads also to an approximation exact to third order (for statistical purposes) for filtered fields. There is, however, an important difference between the two mentioned contributions to the x , because the part due to the tidal field remains small for any power spectra while that due to the averaging process grows indefinitely with increasing power on small scale. This implies, as we shall find, that for power spectra with too much power on small scale the procedure presented here fails even for values of σ smaller than one. However, even in these cases the results are exact to third order, but many orders contribute for finite but small σ . So, the steeper the power spectra the smaller the value of σ at which the approximation start failing. This happens, however, only for very steep power spectra not

very relevant to cosmology. To deal with these cases we should use expression (1) with the mean values of x , not simply the CZA.

3. PDF of the filtered field

The effect of filtering on the field of density fluctuations have been treated exactly by Bernardeau (1994). In this treatment one may, in principle, obtaining (previously to caustic formation) all moments of the PDF of the field of density fluctuation filtered in some scale as a series of powers of σ and sometimes (when the corresponding Hedgeworth expansion converges) to reconstruct the PDF. However, even when this works it is complicated. The procedure described here to deal with this problem, which is basically that in Betancort-Rijo (1991) is more immediate, allowing the direct evaluation of the PDF even for values of σ well above unity. Furthermore, it provides a clearer visualization of the effect of filtering. We now describe the basic idea of our procedure using the CZA, although the procedure in itself is independent of the approximation used for the evolution.

Consider certain geometrical body, B_1 , centered at some point within the initial field. By assigning to this point the mean value of the density fluctuation within B_1 and repeating the process for each point, we generate a new field which we call the field filtered within B_1 . We may now obtain the linear peculiar velocity field corresponding to the growing mode of this field. Let $\lambda_i(B_1)$ be the proper values of the corresponding linear (linearly calculated) deformation tensor, and let B_2 be the body which is the result of transforming B_1 with this deformation tensor. The CZA, in the present context, must be written in the form:

$$1 + \delta(B_2) = \left(\prod_{i=1}^3 (1 - r_i(\vec{\lambda}(B_1))\lambda_i(B_1)) \right)^{-1}. \quad (2)$$

where δ is the present value of the density fluctuations filtered with B_2 at the point under

consideration. In our problem, the filtering body B_2 is a sphere of radius r and B_1 is an ellipsoid with semiaxis $r(1 - r_i(\vec{\lambda})\lambda_i)^{-1}$. Hereafter, we write simply λ_i instead of $\lambda_i(B_1)$. It must be noted that in the actual evolution the present sphere does not transform (when evolved back) exactly into an ellipsoid. The real surface has the same quadrupolar component as the mentioned ellipsoid, but contains smaller wiggles. However, for the present problem those wiggles are completely irrelevant.

Equation (2) gives the best one-point expression for δ as given by the CZA approximation. By one-point expression we mean that the value of δ at a given point (the mean value within a B_2 centered at that point) is expressed only in terms of the λ_i at that point (the mean value within B_1). To go beyond this approximation, we need to obtain the values of the λ_i at different points and integrate over them, or, what is the same, to include the $\mathbf{x}$ variables described in the previous section.

It should be noted that in equation (2) the λ_i depend on B_1 , which, in turn, depends on the λ_i (for a given B_2). It is possible to show that, as long as caustics do not form, for every point there is a unique ellipsoid (with certain axis and orientation) such that the local deformation tensor of the field filtered with the ellipsoid transforms that same ellipsoid into a sphere of radius r (B_2 in general). This prescription uniquely determines the values of the λ_i (hence, δ) at any given point. These unique λ_i (for a given B_2) we call λ'_i . We must now obtain the PDF for the λ'_i at a randomly chosen point in the initial field. With this PDF and equation (2), the derivation of the PDF for δ is a standard problem.

For a field filtered with a given body (the same at every point; usually a sphere of given radius), the PDF for the λ_i is given by (Doroshkevich 1970):

$$P(\lambda_1, \lambda_2, \lambda_3, \sigma) = \frac{5^{5/2}27}{8\pi\sigma^6}(\lambda_1 - \lambda_2)(\lambda_1 - \lambda_3)(\lambda_2 - \lambda_3) \\ \times \exp \left[\frac{-1}{\sigma^2} \left(3(\lambda_1 + \lambda_2 + \lambda_3)^2 - \frac{15}{2}(\lambda_1\lambda_2 + \lambda_2\lambda_3 + \lambda_1\lambda_3) \right) \right];$$

$$\sigma^2 \equiv \langle \delta_L^2 \rangle = \langle (\lambda_1 + \lambda_2 + \lambda_3)^2 \rangle; \quad \lambda_3 < \lambda_2 < \lambda_1. \quad (3)$$

σ^2 is the variance of the initial field of density fluctuations, δ_L , filtered with the given body and extrapolated linearly to the present time. The trace of the deformation tensor extrapolated linearly to the present time (is given the tensor used in the Zel'dovich approximation) by δ_L . Betancort-Rijo (1991) and Padmanabhan & Subramanian (1993) took the PDF for the λ'_i to be simply proportional to the PDF for the λ_i (with $\lambda_i = \lambda'_i$) for the field filtered with the ellipsoid that will contract to a sphere of radius r ; that is, with an ellipsoid with semiaxis $r(1 - r_i(\vec{\lambda}')\lambda'_i)^{-1}$. Note that now the filtering body changes from one point to another, so σ is no longer a constant but a function of the λ'_i .

$$\sigma^2(\vec{\lambda}', r) = \frac{1}{(2\pi)^3} \int W(F)^2 P(K) d^3 \vec{K};$$

$$W(F) = \frac{3[\sin F - F \cos F]}{F^3} \quad (4)$$

with $F = \sqrt{\sum_i K_i^2 E_i^2} r$; $E_i = (1 - r_i(\vec{\lambda}')\lambda'_i)^{-1}$, $K = |\vec{K}|$; $P(K) \equiv$ power spectra.

In practice, in the papers mentioned and, to some extent, here, instead of the full equation (4) a more convenient approximation is used. Since σ depends mainly on the volume of the filtering body (this is exactly so for white-noise power spectra), only the dependence on this quantity is considered. We then have:

$$\sigma^2(\vec{\lambda}', r) = \sigma^2(\vec{0}, r(\prod_{i=1}^3 (1 - r_i(\vec{\lambda}')\lambda'_i)^{-1/3})) \equiv \sigma^2(r(\prod_{i=1}^3 (1 - r_i(\vec{\lambda}')\lambda'_i)^{-1/3})) = \sigma^2(r(1 + \delta)^{1/3}), \quad (5)$$

where $\sigma^2(r)$ is the variance of the linear field filtered with a sphere of radius r . However, if we want to obtain results which are exact to the fourth order, as we shall do in some cases, the small shape dependence of $\sigma(\vec{\lambda}', r)$ must be taken into account. In these cases, we use the following excellent approximation to the actual shape dependence of σ :

$$\sigma^2(\vec{\lambda}, r) = \sigma^2(r(1 + \delta)^{1/3}) Z^2(\vec{\lambda})$$

$$Z(\vec{\lambda}) \approx \exp \left[-\frac{B}{2} \sum_{i < j} \left(\ln \left(\frac{1 - \lambda_i}{1 - \lambda_j} \right)^2 \right) \right] \quad (6)$$

To compute the constant B in the exponential we may use the exact form of Z in the case when two of the λ are equal. In this case and for power law power spectra (with index n) expression (5) reduces to:

$$Z(P, n) = \frac{P^{-\frac{2n}{3}} \int_0^\infty \int_0^\infty (u^2 + P^2 s^2)^{\frac{n}{2}} (W(\sqrt{u^2 + s^2}))^2 s \, ds \, du}{\int_0^\infty W^2(u) u^{2+n} du}$$

$$P \equiv \frac{1 - \lambda_3}{1 - \lambda_1}; \quad \lambda_1 = \lambda_2 \neq \lambda_3 \quad (7)$$

We find: $B(n = -2) = 0.0486$; $B(n = -1) = 0.0499$; $B(n = 0) = 0$.

The approximation used in the mentioned works (which use expression (5)) may be expressed in the form:

$$P'(\vec{\lambda}', \sigma) = C P(\vec{\lambda}, \sigma(r(1 + \delta(\vec{\lambda}))^{1/3}))|_{\vec{\lambda}=\vec{\lambda}'}, \quad (8)$$

where P' is the PDF of the λ'_i ; $\delta(\vec{\lambda})$ is given by equation (2) (with $x=u=0$); σ stands for $\sigma(r)$, and C is a normalization constant. This constant has to be introduced because, unlike P , the integral of P' over all $\vec{\lambda}'$ space is not equal to unity, due to the dependence of σ on $\vec{\lambda}$.

We now show why equation (8) is incorrect and give the correct one. To this end we consider an approximation in which δ depends only on $\delta_L (\equiv \lambda_1 + \lambda_2 + \lambda_3)$. We do this because the relevant fact at this point is basically the same with one and with three variables, but it is much easier to understand in the former case.

For definiteness, let us assume that:

$$1 + \delta = \left(1 - \frac{\delta_L}{3}\right)^{-3}, \quad (9)$$

an approximation that has been used in Betancort-Rijo (1991).

For Gaussian initial conditions, δ_L , which corresponds to a fixed filtering scale, follows a normal distribution. To obtain the PDF for δ'_L (the value of δ_L within the sphere of radius $r/(1 - \frac{\delta_L}{3})$ that have contracted at present to the sphere of radius r) that, by means of equation (9), allows us to obtain the PDF for δ , we proceed as follows. To say that at a given point δ'_L takes the value δ_L^0 means that at this point the linear value of δ filtered with a sphere of radius $r/(1 - \delta_L/3)$ is just δ_L^0 . The accumulated probability, $P'(\delta'_L \geq \delta_L^0)$, that at a point chosen at random in the initial field δ'_L is larger than or equal to some given value may be obtained immediately. To this end we only need noting that if a sphere of radius $r/(1 - \delta'_L/3)$ (with $\delta'_L \geq \delta_L^0$) have to contract to a sphere of radius r and shell crossing does not happen, the sphere of radius $r/(1 - \delta_L^0/3)$ will have to contract to a radius smaller than r . So, the value of δ_L within the sphere with radius $R/(1 - \delta_L^0/3)$ must be larger than δ_L^0 , since if the value of δ_L within this sphere were just δ_L^0 it would contract to a sphere of radius r . Thus, the accumulated probability under consideration must be equal to the probability that the value of δ_L within a sphere of radius $r/(1 - \delta_L^0/3)$ be larger than or equal to δ_L^0 . So, we have

$$P'(\delta'_L \geq \delta_L^0) = \frac{1}{2} \text{erfc} \left(\frac{\delta_L^0}{\sqrt{2}\sigma(\delta_L^0)} \right)$$

$$\sigma(\delta_L^0) = \sigma \left(r \left(1 - \frac{\delta_L^0}{3} \right)^{-1} \right) \quad (10)$$

where erfc stands for the error complementary function. To obtain the probability distribution for the values of δ'_L (the value of δ_L within the sphere that will contract to a sphere of radius r) at a randomly chosen point we only need deriving the cumulative probability with respect to δ_L^0 .

$$\begin{aligned}
P'(\delta'_L) &= -\frac{d}{d\delta_L^0} P'(\delta_L > \delta_L^0)|_{\delta_L^0=\delta'_L}; \\
P'(\delta'_L)d\delta'_L &= \frac{1}{\sqrt{2\pi}} e^{-\delta'^2_L/(2\sigma^2(\delta'_L))} d\left(\frac{\delta'_L}{\sigma(\delta'_L)}\right).
\end{aligned} \tag{11}$$

Note that since σ depends on δ'_L we cannot take it out of the differential. So we cannot write:

$$\begin{aligned}
P'(\delta'_L) &= C P(\delta_L, \sigma(\delta_L))|_{\delta_L=\delta'_L} \\
&= C \frac{e^{-\delta'^2_L/(2\sigma^2(\delta'_L))}}{\sqrt{2\pi}\sigma(\delta'_L)},
\end{aligned}$$

contrary to the assumption in equation (8). With these considerations in mind, it is easy to obtain the correct distribution for the λ'_i . We need only to generalize the above argument to three dimensions. The key argument is that the variables $\lambda'_i/\sigma(\vec{\lambda}')$ must follow the distribution followed by the variable λ_i/σ for fixed filtering (when σ is a constant). So we have:

$$\begin{aligned}
P'(\vec{\lambda}', \sigma) &= \left(P(\vec{\lambda}, \sigma(\vec{\lambda})) \sigma^3(\vec{\lambda}) \left| \frac{\partial \frac{\lambda_i}{\sigma(\vec{\lambda})}}{\partial \lambda_j} \right| \right) |_{\vec{\lambda}=\vec{\lambda}'}; \\
\sigma(\vec{\lambda}) &\equiv \sigma(\vec{\lambda}, r).
\end{aligned} \tag{12}$$

The last factor in this expression is the Jacobian of the transformation from the variables $\lambda_i/\sigma(\lambda_i)$ to the variables λ_i . Note that the P' obtained by means of this expression is automatically normalized.

Equation (12) is exact for any initial conditions (provided that the λ_i determine the evolution), although we shall use it only for Gaussian initial conditions; that is, with P given by equation (3).

P' is the PDF of the λ'_i (the λ_i associated with the unique ellipsoid that will contract to the sphere of radius r centered at the given point) for a point chosen at random in the initial field; that is, in Lagrangian coordinates. However, we want to obtain the PDF for δ at a point chosen at random in the present “physical” coordinates (i.e. Eulerian coordinates). So, we must first obtain the PDF for the λ'_i in Eulerian coordinates. To this end, we only need to multiply P' by the Jacobian of the transformation from Lagrangian to Eulerian coordinates. For a non-filtered field this Jacobian is simply

$$\prod_{i=1}^3 (1 - r_i(\vec{\lambda})\lambda_i). \quad (13)$$

However, for filtered fields this expression is not correct. This can be appreciated by noting that the resulting distribution for the λ'_i (the result of multiplying P' by the Jacobian in question) is not normalized. For non-filtered fields, on the other hand, it is easy to check that the Jacobian given by equation (13) leads to a normalized PDF. The correct Jacobian must exhibit the following properties: it has to be zero at points where δ goes to infinity; it must reduce to expression (13) in the limit of small power on scales below r ; it must render a normalized PDF with vanishing first moment for any power spectra. The following expression meets these requirements and is exact at least to first order

$$J(\lambda) = \prod_i (1 - (1 - \gamma)r_i(\vec{\lambda})\lambda_i), \quad (14)$$

where γ is certain spectral constant that tends to zero in the limit of small power on scales below r . We cannot say a priori whether this expression is exact to higher orders. However, it seems to be exact to second order (within the precision of our computations) and it gives excellent results to all orders. It must be noted that expression (14) is not the value of the Jacobian at a point with the given λ_i . This would be so if the evolution were determined solely by these quantities. However, we know that this is not strictly true, so

expression (14) should represent the mean value of the Jacobian over all points with a given value of the λ_i . To obtain γ we have considered a fluctuation with spherical symmetry and the evolution of the region such that the mean density fluctuation within a sphere centered at points belonging to it is above some threshold. Using the spherical model with the mean density profile, we obtained the contraction of this region and by comparison with expression (14) we find γ . However, this derivation is lengthy and although it might be interesting in itself, it is not essential to the present purposes, since we may obtain that constant using any relevant analytical result available (i.e., the leading order value of S_3) and comparing it with the one obtained using expression (14). In either case, we find:

$$\gamma = \frac{1}{6} \frac{d[\ln(\sigma^2(r))]}{d[\ln(r)]} \quad (15)$$

4. Results and conclusions

To check the accuracy of the approximation developed in this article, we shall present some explicit calculations and compare them with the results of the corresponding available numerical simulation and with exact results obtained with the perturbation theory. We shall consider Gaussian initial conditions with power spectra of the form $P(k) = Ck^n$.

In these cases, equation (5) takes the form:

$$\begin{aligned} \sigma(\vec{\lambda}, r) &= \sigma_0(1 + \delta)^{-\alpha/2}; \\ \sigma_0 &\equiv \sigma(r); \alpha \equiv \frac{n+3}{3}, \end{aligned} \quad (16)$$

with δ given by the CZA

$$(1 + \delta)^{-1} = \prod_i (1 - r_i(\lambda) \lambda_i) \quad (17)$$

To derive exact results we use for $r_i(\lambda)$ its exact expression (Betancort-Rijo & López-Corredoira 2000). To derive result for finite values, however, we find it much more convenient to use the following compact approximation

$$r_i(\lambda) = g\left(\frac{3}{2}(\lambda_J + \lambda_K), \Omega\right)$$

$$g(x, \Omega) = \begin{cases} \frac{3}{x} \left(1 - \frac{x\Omega^{2/63}}{1.66}\right)^{0.5533} & ; x \leq 1.584 \\ 1.55 & ; x > 1.584 \end{cases} \quad (18)$$

Using this in expression (12), with $P(\vec{\lambda}, \sigma)$ given by (3) and multiplying it by the Eulerian–Lagrangian Jacobian, $J(\vec{\lambda}, \alpha)$, (expression (14)), we obtain the PDF for the λ'_i corresponding to a point chosen at random in Eulerian coordinates, which we represent by $G(\vec{\lambda}', \sigma_0)$:

$$G(\vec{\lambda}', \sigma_0) = \left(\left(P(\vec{\lambda}, \sigma(\vec{\lambda})) \sigma^3(\vec{\lambda}) \left| \frac{\partial \frac{\lambda_i}{\sigma(\vec{\lambda})}}{\partial \lambda_j} \right| \right)_{\vec{\lambda}=\vec{\lambda}'} \right) J(\vec{\lambda}', \alpha). \quad (19)$$

The moments of the PDF of the present value of the density fluctuation top-hat filtered on scale r are then given by:

$$\langle \delta^n \rangle(\sigma_0) = \int_{-\infty}^{\infty} d\lambda'_1 \int_{-\infty}^{\lambda'_1} d\lambda'_2 \int_{-\infty}^{\lambda'_2} d\lambda'_3 G(\vec{\lambda}', \sigma_0) \delta^n, \quad (20)$$

where δ is given by equations (17) with $\vec{\lambda} = \vec{\lambda}'$, provided that each of the factors in (17) is positive and δ is smaller than 170. If this expression leads to a value of δ larger than this, we use the following expression:

$$\delta = 170 \left| \frac{\delta_L}{1.589} \right|^3.$$

This last prescription represents an attempt to deal with caustic formation in a simple

manner, so that equation (23) gives sensible results for any value of σ . However, it is not relevant for the results given here.

The PDF for δ , $P(\delta, \sigma_0)$, may be obtained in the very process of computing any of the moments. To this end, we add up all the values of $G(\vec{\lambda}, \alpha)$ (multiplied by the corresponding volume of the integration step) that are found during the integration to lie between δ and $\delta + \Delta\delta$ (for a suitably chosen $\Delta\delta$, depending on δ) and divide the result by $\Delta\delta$.

The values of δ given by (17) for very under dense regions is not correct this imply $\langle\delta\rangle \neq 0$. However, this problem disappears if we set $P(\delta) = 0$ for $\delta < -0.82$.

The values of S_3 and S_4 given by

$$S_3 \equiv \frac{\langle\delta^3\rangle}{\langle\delta^2\rangle^2}; \quad S_4 \equiv \frac{\langle\delta^4\rangle - 3\langle\delta^2\rangle^2}{\langle\delta^2\rangle^3}$$

in the low- σ limit, may be computed analytically. The exact value of $S_3(\sigma = 0) = (34/7 - (n + 3))$ is obtained, for the power spectra under consideration, when expression (17) for δ is used; the value $4 - (n + 3)$ is found when the r_i are set equal to one. For a non-filtered field we have computed $S_4(\sigma = 0)$ analytically both for the CZA (equation (17)) and for the Zel'dovich approximation. Note that the non-filtered case may be treated formally as a filtered field with negligible power below the filtering scale, so that we may use expression (30) with σ independent of λ and $\gamma = 0$. For power law power spectra this corresponds to the limit $n = -3$, but it corresponds more generally to any power spectra having the mentioned properties. The correct values (60712/1323; 279/9; Bernardeau 1994a) are found. For the evolution of these amplitudes and the variance in the small σ limit we find

$$\langle\delta^2\rangle = \sigma^2 + 1.81\sigma^4 + 3.8\sigma^6$$

$$S_3 = 4.86 + 9.815\sigma^2 + 48\sigma^4$$

$$S_4 = 45.89 + 288.8\sigma^2 + 2072\sigma^4 \quad (21)$$

This results are exact to next leading order for S_3 (in principle only for $n = -2.7$ but the n dependence is negligible) and almost exact to the same order for $\langle\delta^2\rangle$, S_4 . As pointed out in section 2, we could use the statistical properties of the x , u variables to obtain these two quantities exactly to the stated order, but the second coefficient in the expansion for $\langle\delta^2\rangle$ will increase by less than a per cent which is about the numerical error for this coefficient. So, it would be an unnecessary complication to use here those additional variables. Furthermore, the purpose of the present procedure is to obtain directly and accurately the PDF for finite values of σ . The question of the maximum order of exactness, although interesting, is only of peripheral relevance to our purposes. These results agree with those exact results given by the perturbation theory by Scoccimarro & Frieman (1995) and the remaining ones are close to those obtained by Fosalba and Gaztañaga (1998) using the spherical model. We see that this model predict well the hierarchical amplitudes but not so well the moments. The above results provide a check to the CZA but not properly to the procedure presented here, since for non-filtered fields expressions (12) and (14) reduce to the standard form. A check to these expressions is provided by the cases of power law power spectra with $n \geq -3$. For $n = -2$ we find:

$$\begin{aligned} \langle\delta^2\rangle &= \sigma^2 + 0.88\sigma^4 + 0.54\sigma^6 \\ S_3 &= 3.86 + 3.12\sigma^2 \\ S_4 &= 27.56 + 58\sigma^2 \end{aligned} \quad (22)$$

Here, to obtain the second coefficient in $\langle\delta^2\rangle$ exactly we have considered the small ($=0.015$) contribution due to the dispersion of the x variables. This is not necessary for the amplitudes. It is also important to consider the shape dependence of σ (see expression

(6)), otherwise we would obtain a coefficient 0.1 too large. For $n = -1$, $n = 1$ we find respectively:

$$\langle \delta^2 \rangle = \sigma^2 + 0.223\sigma^4$$

$$S_3 = 2.86 + 0.267\sigma^2$$

$$S_4 = 13.89 - 2.583\sigma^2$$

$$\langle \delta^2 \rangle = \sigma^2 - 0.173\sigma^4$$

$$S_3 = 0.86 + 0.8\sigma^2$$

$$S_4 = 0.556 - 17.8\sigma^2 \tag{23}$$

For $n \geq -1.5$, S_3 and S_4 are practically independent of σ . For example, for $n = -1.3$ the change of these quantities up to $\sigma = 1.52$ is less than a 4%. For smaller values of n , however, the evolution of this quantities is more apparent. For $n = -2$ we find (table 1) that S_3 and S_4 increase considerably from $\sigma = 0$ to $\sigma \approx 0.5$ and more steeply for higher values of σ . In figure 1 we present the PDF for δ corresponding to $P(k) \propto k^{-1}$ and $\sigma = 0.3$; 0.5 . Comparing this with figure 7 of Bernardeau & Koffman (1995), we find them to be almost identical. The PDFs of Bernadeau & Koffman were obtained under the assumption that the S_p parameters (the p -th cumulants divided by $\sigma^{2(p-1)}$) are independent of σ . We have seen that this is the case for $n = -1$, at least for S_3 and S_4 . So, the agreement between both pairs of PDFs is to be expected. The evolution of $\langle \delta^2 \rangle$ agrees very well with the result by Lokas et al. (1996).

In figure 2 the PDFs corresponding to $\sigma = 0.92$; 1.52 , $n = -1.3$ are represented. These should be compared with figure 10 of Bernardeau & Kofman (1995). The agreement is

very good, our results being always within the numerical errors of the results given in that article.

It should be noted that the mentioned numerical results assume a cold dark matter (CDM) while we have used a power-law power spectrum with the same local logarithmic derivative. Using the whole CDM power spectrum in our computations would represent only a small complication. If we have not done so, it is because this implies only a slight improvement in the results that is completely unobservable given the precision of the simulations.

The procedure described here for obtaining the PDF for δ is in principle valid only for top-hat filtered fields. However, the basic lines of reasoning followed here are entirely translatable to another smoothing procedures.

For a given smoothing (in the present field), we must determine the smoothing in the initial field rendering the minimum dispersion for the smoothed present field. That is, we must find a smoothing procedure that, when applied to the initial field at a given point, renders a set of λ'_i (extrapolated linearly to the present time) that predict the mean value of δ at the transformed point, given by an expression similar to (17), and so that the dispersion in δ for given λ'_i s is a minimum. This is what we have implicitly done here for the spherical top-hat smoothing, and we have found that it corresponds to an initial ellipsoidal top-hat smoothing.

The solution to this problem for Gaussian smoothing is not simple, although one could guess that a triaxial Gaussian, with the axes determined as in the top-hat case, should not be a bad approximation. Within this simple approximation and for a power-law spectrum the PDF for δ must be the same as for the top-hat case with the same σ . For more general power spectra, we cannot decide a priori whether the value of the logarithmic derivative must be taken on the scale where the top-hat has the same volume as the Gaussian or on

the scale where it has the same σ .

As an example we discuss one of the cases represented in figure 4 of Kofman et al. (1994), where the Gaussian smoothing radius is $R_s = 6h^{-1}\text{Mpc}$; $\sigma = 0.55$. If we matched the volumes, the corresponding top-hat radius is $9.38 h^{-1}\text{Mpc}$, so that for a CDM power spectrum $n \approx -1$. The PDF corresponding to a top-hat filtering with $\sigma = 0.55$; $n = -1$ is represented in figure 3 and shows very good agreement with the numerical results in the last reference. If we matched the σ 's, the corresponding top-hat smoothing scale imply $n = -1.1$. The corresponding PDF is almost indistinguishable from the former.

We have seen that to obtain the PDF for δ we must integrate over all values of λ subject to the condition $\delta(\lambda) = \delta$. If we considered, however, an approximation where δ depended on the λ_i only through $\delta_L = \lambda_1 + \lambda_2 + \lambda_3$, the computation of $P(\delta)$ would be greatly simplified. We have already seen that such an approximation cannot be entirely correct, but leads to a simple analytical expression that displays, in a clear manner, the effect of smoothing. Taking for $\delta(\delta_L)$ that corresponding to spherical collapse and using the fact that the distribution for δ_L is simply a Gaussian, equation (11) and (14) immediately gives for $P(\delta)$:

$$\begin{aligned}
 P(\delta) &= \frac{1}{\sqrt{2\pi}} \frac{e^{-\frac{\delta_L^2(\delta)}{2\sigma^2(\delta)}}}{\sigma(\delta)(1+\delta)} \left((1+\delta)^{-0.602} + \frac{\alpha}{2} \delta_L(\delta) \right) \\
 &\quad \times \left(1 - \left(1 - \frac{\alpha}{2} \right) \frac{\delta_L(\delta)}{1.66} \right)^{1.66}; \\
 \sigma(\delta) &= \sigma_0 (1+\delta)^{-\alpha/2}; \quad \alpha \equiv \frac{n+3}{3} \\
 \delta_L(\delta) &= 1.66(1 - (1+\delta)^{-0.6024}).
 \end{aligned} \tag{24}$$

Where the last equation represents a very accurate approximation to the spherical model that we propose. However, expression (24) represents only a rough approximation to

$P(\delta)$. For $\sigma = 1.52$ the value of $P(\delta)$ given by (24) may differ from the real value by a factor greater than three. Furthermore, the value of $S_3(\sigma = 0)$ given by (24) is $37/7 - (n + 3)$, which is $3/7$ too large.

We may conclude that for $\delta \in (-0.9, 10)$ and σ as large as 1.52, the $P(\delta)$ computed as indicated here is, so far as we have been able to check, in excellent agreement with the results of numerical simulations. This agreement is particularly meaningful since our derivation of $P(\delta)$ involves neither free parameters nor modeling freedom (except for some minute refinements non-essential for the results given here). Equation (12) is exact and is valid for any field. Equation (14) for the Jacobian of the Eulerian–Lagrangian transformation in a filtered field is also exact (at least to first order in λ_i) and, although derived for a Gaussian field, it seems to be valid for any field.

REFERENCES

- Bernardeau, F. 1994a, ApJ, 433, 1
- Bernardeau, F. 1994b, ApJ, 427, 51
- Bernardeau, F., & Kofman L. 1995, ApJ, 443, 479
- Betancort-Rijo, J. 1991, MNRAS, 251, 399
- Betancort-Rijo J., López-Corredoira M., 2000, ApJ 534, L117
- Betancort-Rijo J., 2001, in preparation
- Bouchet, F. R., Colombi, S., Hivon, E., & Juszkiewicz. R. 1995, A&A, 296, 575
- Buchert, T. 1989, A&A, 223, 9
- Doroshkevich, A. G. 1970, Astrofizica (VI) 6, 4, 581
- Fosalba P, Gaztañaga E., 1998, MNRAS 301, 503
- Hoffman, V. 1987, ApJ, 318, L7
- Kofman, L., Bertschinger, E., Gelb, J., Nusser, A., & Dekel, A. 1994, ApJ, 420, 44
- Padmanabhan, T., & Subramanian K. 1993, ApJ, 410, 482
- Scoccimarro R., Frieman J., 1995, ApJ 473, 620
- Tokas E. L., Juszkiewicz R., Bouchet F. R., Hivon E., 1996, ApJ 467, 1
- Trimble, V. 1987, ARA&A, 25, 425
- Zel'dovich, Y. B. 1970, A&A, 5, 20

FIGURE CAPTIONS

Fig. 1: Probability distribution function (PDF) for $\sigma = 0.3$ and $\sigma = 0.5$; $n = -1$.

Fig. 2: Probability distribution function (PDF) for $\sigma = 0.92$ and $\sigma = 1.52$; $n = -1.3$.

Fig. 3: Probability distribution function (PDF) for $\sigma = 0.55$; $n = -1$.

PDF

$n=-1$





Table 1: $\langle \delta^2 \rangle$, S_3 and S_4 as a function of σ and n .

σ	n	$\langle \delta^2 \rangle$	S_3	S_4
0.3	-2	0.0950	4.06	32.4
0.3	-1	0.0889	2.91	13.4
0.5	-2	0.290	4.51	44.5
0.5	-1	0.243	3.03	14.9
0.75	-2	0.670	4.54	43.0
0.75	-1	0.527	2.96	16.6
1.0	-2	1.51	5.4	77
1.0	-1	0.85	2.84	17.0
1.52	-2	5.1	8.5	124
1.52	-1	1.43	2.68	16.5

PDF

$n=-1.3$



PDF

$n=-1$

